



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07K 7/40, G01N 33/532, 33/533	A1	(11) International Publication Number: WO 91/16344 (43) International Publication Date: 31 October 1991 (31.10.91)
(21) International Application Number: PCT/US91/02212 (22) International Filing Date: 4 April 1991 (04.04.91) (30) Priority data: 512,272 20 April 1990 (20.04.90) US (71) Applicant: GEORGETOWN UNIVERSITY [US/US]; Ryan Administration Building, 37th & O Streets, N.W., Washington, DC 20057 (US). (72) Inventors: BREDEHORST, Reinhard ; 2617 42nd Street, N.W., Washington, DC 20007 (US). LIGLER, Frances, S. ; 12400 Bealle Mount Road, Potomac, MD 20854 (US). KUSTERBECK, Anne, W. ; 6428 Quincy Place, Falls Church, VA 22040 (US). WEMHOFF, Gregory, A. ; 10323 7th Regiment Drive, #203, Manassas, VA 22110 (US). VOGEL, Carl-Wilhelm ; 4100 Massachussetts Avenue, N.W., Washington, DC 20007 (US).		(74) Agents: OBLON, Norman, F. et al.; Oblon, Spivak, McClelland, Maier & Neustadt, Crystal Square Five, Fourth Floor, 1755 South Jefferson Davis Highway, Arlington, VA 22202 (US). (81) Designated States: AT (European patent), BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i>
(54) Title: LIGAND-LABEL CONJUGATES WHICH CONTAIN POLYOXOANIONS OF SULFUR OR PHOSPHORUS (57) Abstract Ligand-label conjugates which are an oligopeptide of 5 to 100 amino acid residues, bonded to a ligand or receptor, which contain a plurality of chemiluminescent or fluorescent labels and a plurality of polyoxoanions of sulfur or phosphorus are useful for immunoassays. Such conjugates are hydrophilic and exhibit very low nonspecific binding, thereby significantly increasing the signal to background ratio in immunoassays.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

DescriptionLigand-Lab 1 Conjugat s Which Contain
Polyoxoanions of Sulfur or PhosphorusTechnical Field

The present invention relates to novel fluorescent-
5 labeled or chemiluminescent-labeled conjugates for use in
specific binding assays for ligands (e.g., antigens and
haptens) or receptors (e.g., antibodies, specific binding
proteins, and cell surface receptors). The invention
10 further relates to intermediate compounds produced in the
synthesis of the novel labeled conjugates, and kits
containing such labeled conjugates and/or intermediates.

Background Art

Presently, several immunochemical methods exist for
the detection of ligands such as haptens, antigens or
15 antibodies. Radioimmunoassay is a widely used method.
This method employs a radioisotope-labeled antigen
(ligand) to compete with the antigen from a test sample
for binding to a specific antibody. U.S. Patent No.
3,709,868 describes such a radioimmunoassay. While by
20 definition, radioimmunoassay is based on the binding of a
specific antibody with an antigen or hapten, radioactive
binding assays which are based on other specific binding
interactions, such as between hormones and their binding
proteins, have also been developed.

25 Several non-isotopic immunoassays have been proposed
to eliminate the disadvantages associated with radioactive
materials. Ligands such as an antigen or hapten and
receptors such as an antibody have been labeled with a
variety of non-radioactive compounds, including
30 chemiluminescent and fluorescent molecules.

Specific examples of useful chemiluminescent labels are disclosed in German OLS No. 2,618,511 and include luminol (3-aminophthalhydrazide or 5-amino-2,3-dihydro-1,4-phthalazinedione) and isoluminol (4-aminophthalhydrazide or 6-amino-2,3-dihydro-1,4-phthalazinedione). The use of N-(4aminobutyl)-N-ethylisoluminol (6-[N-(4-aminobutyl)-Nethylamino]-2,3-dihydrophthalazine-1,4-dione) as a chemiluminescent label is reported in Simpson et al., Nature, Vol. 279, p. 646 (1979). The preparation of chemiluminescent phthalhydrazide labeled ligands is described in U.S. Patent Application Serial No. 927,621, filed July 24, 1978, entitled "Chemiluminescent Phthalhydrazide Labelled Conjugates." U.S. Patent 4,331,808 discloses labeling ligands with chemiluminescent naphthalene-1,2-dicarboxylic acid hydrazide using straight chain alkyl groups as spacers between the label and the ligand.

While these chemiluminescent compounds have been shown to be suitable labels, they are hydrophobic and, thereby, increase the hydrophobicity of the ligand complex. For example, column chromatographic purification of synthesized thyroxine-naphthalene-1,2-dicarboxylic acid hydrazide conjugates requires the use of organic solvents (U.S. Patent No. 4,331,808). A documented disadvantage of an increased hydrophobicity is an increase in nonspecific binding effects. This results in a decreased signal to background ratio of immunoassays.

U.S. Patent No. 4,645,646 describes the use of hydrophilic chain-like polymers with recurring functional groups (e.g., proteins) as carriers of multiple chemiluminescent luminol molecules to improve the sensitivity of luminescence immunoassays. However, these

conjugates exhibit non-specific binding properties. In fact, some of the disclosed conjugates exhibit such marked non-specific binding properties to preclude their use in immunoassays (U.S. Patent No. 4,645,646). These examples
5 demonstrate that currently available techniques for coupling chemiluminescent labels to ligands pose an inherent problem. The hydrophobic properties of the chemiluminescent labels decrease the water solubility of the ligands to an extent that the sensitivity of the
10 immunoassay is significantly reduced due to high non-specific background binding.

Labeling of ligands with fluorescent molecules poses very similar problems. Evrain et al., Steroids, vol. 35, 611-619 (1980) describe the synthesis of three
15 fluorescein-labeled derivatives of testosterone using either cysteamine, or 1,3-diaminopropane, or 1,7-diaminoheptane as spacer between the fluorophore and the ligand. All derivatives proved to be highly hydrophobic. For example, analysis of the testosterone
20 fluorescein conjugates by thin-layer chromatography on silica gel required the use of a combination of benzene:ethyl acetate:acetone (1:8:1) or chloroform:ethanol (7:3) as the solvent system. The hydrophobic nature of this compound is typical of
25 fluorescein-labeled molecules and generally leads to high background readings in immunoassays.

U.S. Patent 4,670,406 describes the use of bifunctional aromatic compounds (e.g., paranitrophenylisocyanate) as rigid coupling compounds for
30 the synthesis of labeled ligands such as fluorescein-labeled digoxin. These rigid coupling reagents are advantageous in that they do not permit the fluorescent

marker to "fold back" onto the ligand, thereby minimizing the possibility of quenching of the fluorescent compound by the ligand. However, as a result of their hydrophobic properties, the rigid coupling reagents further contribute
5 to the loss of the water-solubility of the ligands upon labeling with fluorophores.

U.S. Patent 4,452,886 describes the synthesis of ligand-containing polymers as carriers of multiple photon emitting (fluorescent) compounds. The disclosed polymers
10 include proteins and synthetic or natural polypeptides having a large number of diamino acids for covalent attachment of photon emitting compounds or polymers of such compounds. Similar approaches have also been described in other publications. U.S. Patent 4,604,364
15 discloses tracer compositions for immunoassays which contain photon emitting compounds coupled to ligands via an intermediate support material such as a protein or polypeptide. U.S. Patent Nos. 4,166,105 and 4,169,137 describe antigen detecting reagents which are prepared by
20 covalently linking fluorescent dye molecules to an appropriate antibody through a polymeric backbone having reactive functional groups along the length of its chain. Polymer backbone molecules reported to be suitable are polyethyleneimines (molecular weight range 1200 to 60,000
25 daltons), polypeptides such as polylysines, polyamides such as nylon-6, and low molecular weight (100 to 10,000 daltons) polymeric carboxylic acids. While such polymers help to reduce the loss of antibody binding activity upon fluorescent labeling, they are not suitable to compensate
30 sufficiently for the hydrophobic properties of fluorescent labels such as fluorescein. As a result, such antigen detecting reagents exhibit significant non-specific background binding, thereby limiting the signal to

background ratio of immunoassays. This is clearly demonstrated by a recent observation, in which antibody, molecules labeled with only three fluorescein residues per antibody exhibited an approximately ten-fold higher non-specific binding than the same antibodies labeled with iodine-125 (personal communication, Lisa Shriver-Lake (1990)).

Specific and Non-specific Binding for DTAF- and ^{125}I -Labeled Goat IgG (Shriver-Lake (1990)).

	Labeled Antigen	Amount of Labeled Antigen Bound to Immobilized anti-Goat IgG (Specific Binding)	Amount of Labeled Antigen Bound to Immobilized Non-immune IgG (Nonspecific Binding)	Ratio of Specific to Nonspecific Binding
20	^{125}I -labeled Goat IgG	447.9 ng	30.4 ng	14.73
25	DTAF-labeled Goat IgG (3.1 DTAF/IGG)	12.97 FU	9.52 FU	1.36

FU: arbitrary fluorescence units; DTAF: 5-(4,6-dichlorotriazinyl)aminofluorescein. The specific and nonspecific binding ratios for the DTAF-labeled goat IgG and the ^{125}I -labeled goat IgG were measured using affinity purified anti-goat IgG (Jackson ImmunoResearch, West Grove, PA) covalently attached to cover slips by N- γ -maleimidobutyryloxy succinimide ester (GMBS) as described in Bhatia et al, Anal. Biochem., vol. 178, pp. 408-413 (1989).

Thus, even a highly hydrophilic polypeptide chain such as an antibody with a molecular weight of approximately 150,000 apparently cannot compensate for the hydrophobic properties of as few as three fluorophore
5 molecules.

Accordingly, there remains a need for molecular carriers which can link fluorescent or chemiluminescent labels to ligands, such as antigens and haptens, or receptors, such as antibodies, specific binding proteins
10 and cell surface receptors, which do not suffer from the above-mentioned drawbacks. In particular, there remains a need for fluorescent-labeled and chemiluminescent-labeled conjugates which possess good water solubility, and for compounds which contain a plurality of fluorescent or
15 chemiluminescent labels or sites for attaching such labels and which may be conveniently covalently bonded to a ligand, such as an antigen, hapten, or a receptor, such as an antibody, etc., to provide a ligand-label conjugate with good water solubility.

20 Disclosure of the Invention

Accordingly, it is an object of the present invention to provide novel conjugates in which

chemiluminescent or fluorescent labels are linked to ligands or receptors.

It is another object of the present invention to provide novel conjugates, in which chemiluminescent or
5 fluorescent labels are linked to ligands and receptors, which are hydrophilic.

It is another object of the present invention to provide novel conjugates, in which chemiluminescent or fluorescent labels are linked to ligands and receptors, in
10 which the quenching of the label by the ligand is minimized.

It is another object of the present invention to provide diagnostic kits which contain such conjugates of ligands or receptors and fluorescent or chemiluminescent
15 labels.

It is another object of the present invention to provide novel labeled compounds in which a plurality of fluorescent or chemiluminescent labels are bonded to a
linker which possesses a unique functional group for
20 bonding a ligand or a receptor.

It is another object of the present invention to provide novel ligand compounds in which a ligand or a receptor is bonded to a linker molecule which possesses a plurality of functional groups for bonding a plurality of
5 fluorescent or chemiluminescent labels.

These and other objects, which will become apparent during the course of the following detailed description have been achieved by ligand-label conjugates in which an oligopeptide is bonded to a ligand or receptor, and in
10 which at least one of the amino acid residues contains a polyoxoanion group of sulfur or phosphorus and a plurality of the amino acid residues are linked to a chemiluminescent or fluorescent label; the intermediate compounds for coupling the ligand and/or labels to such
15 conjugates; and kits containing such conjugates.

Brief Description of the Drawings

A more complete appreciation of the invention and many of the attendant advantages thereof will be readily obtained as the same become better understood by reference
20 to the following detailed description when considered in connection with the accompanying drawings, wherein:

Figure 1 illustrates the fluorescence lifetime of the DNP-insulin A-chain-fluorescein conjugate. Shown are the frequency-dependent phase (\square) and modulation (\blacksquare) data, together with the best one (- - -) and two (---) component
5 fits. The precision of the data is approximately ten-fold better (phase: 0.2 degrees; modulation: 0.002) than suggested by the size of the data points;

Figure 2 illustrates the relationship between the fluorescence intensity and concentration of the DNP-
10 insulin A-chain-fluorescein conjugate and the DNP-lysine-fluorescein conjugate; and


Figure 3 illustrates the binding of different DNP-conjugates to immobilized monoclonal anti-DNP antibody 51.

Best Mode for Carrying Out the Invention

15 In one aspect, the present invention relates to conjugates in which a natural or synthetic polypeptide oligomer is bonded to a ligand or receptor, and in which at least one of the amino acid residues contains a
polyoxoanion of sulfur or phosphorus and a plurality of
20 the amino acid residues are linked to a chemiluminescent or fluorescent label. The polypeptide backbone of the molecule may suitably contain 5 to 100 amino acid

residues, preferably 10 to 50 amino acid residues, most preferably 15 to 25 amino acid residues.

The present conjugates may thus be represented by the general formula (I):




wherein n is an integer of from 5 to 100, preferably 10 to 50, most preferably 15 to 30, and $(aa_1) \dots (aa_n)$ represent amino acid residues, and wherein at least one of the amino acid residues contains a polyoxoanion of sulfur
10 or phosphorus, a plurality of the amino acid residues are linked to a fluorescent or chemiluminescent label, and one of the amino acid residues is bonded to a receptor or ligand.

The present conjugates are easily prepared from
15 synthetic or natural polypeptide oligomers which contain three types of functional groups: (i) a unique first functional group for bonding a ligand or receptor; (ii) at least one of a second functional group which may be derivatized to contain a polyoxoanion of phosphorus or
20 sulfur; and (iii) a plurality of a third functional group which can be linked to a chemiluminescent or fluorescent label. Alternatively, the starting oligopeptide may be

one which already contains at least one polyoxoanion, such as insulin A-chain in the tetra-S-sulfonate form and thus, is not required to possess the second type of functional group described above.

5 The unique functional group for bonding the ligand is preferably located at one end of the polypeptide oligomer chain. Accordingly, it is preferred that this functional group be either a NH_2 - group or a $-\text{CO}_2\text{H}$ group, since these reactive groups are the natural terminators of
10 polypeptide chains.

In this case, the conjugate may have either formula (II) or (III):



15 in which the definitions of formula (I) apply and L is a ligand or receptor.

It is particularly important that the functional group for bonding the ligand be a unique group and that only one ligand is attached to the polypeptide backbone.
20 This is Of particular importance for the detection of

small molecular weight ligands in competitive immunoassays. Since the signal in such assays is usually generated by the release of labeled ligand from the antibody, increased sensitivity is achieved if the ligand
5 is bound by only one antibody binding (Fab') site. If two or more ligands were included in the labeled conjugate, one ligand could be displaced while another remained bound to, for example, an antibody-coated solid support.

The term ligand, as used herein, refers to a
10 molecule such as an antigen or hapten which binds to a corresponding receptor such as an antibody. Generally, such ligands include, for example, drugs, hormones, proteins, vitamins, and infectious agents. The term receptor, as used herein, includes, for example,
15 antibodies, specific binding proteins, and cell surface receptors. As discussed above, the present conjugates containing ligands and receptors are useful reagents for quantitative and qualitative measurements of the same ligand or receptor in homogeneous and heterogeneous
20 immunoassay systems.

Representative of receptors are antibodies in general, particularly those of the IgG, IgE, IgM and IgA classes, for example hepatitis B antibodies; and representative of ligands are antigenic proteins such as

-13-

insulin, chorionic gonadotropin (e.g., HCG),
carcinoembryonic antigen (CEA), myoglobin, hemoglobin,
follicle stimulating hormone, human growth hormone,
thyroid stimulating hormone (TSH), human placental
5 lactogen, thyroxine binding globulin (TBG), intrinsic
factor, transcobalamin, enzymes such as alkaline
phosphatase and lactic dehydrogenase, hepatitis-associated
antigens such as hepatitis B surface antigen (HB_sAg),
hepatitis r antigen (HB_eAg) and hepatitis core antigen
10 (HB_cAg) and any antigens derived from HIV such as, e.g.,
gp120. Representative of polypeptide ligands are
angiotensin I and II, C-peptide, oxytocin, vasopressin,
neurophysin, gastrin, secretin, and glucagon.


Since, as peptides, ligands of this general category
15 comprise numerous available carboxylic acid and amino
groups, coupling to either the terminal -NH₂ or -CO₂H group
of the oligopeptide backbone can proceed according to
conventional peptide condensation reactions such the
carbodiimide reaction, the mixed anhydride reaction, and
20 so forth, or by the use of conventional bifunctional
reagents capable of coupling carboxylic acid or amino
functions to either the terminal -NH₂ or -CO₂H group of the
oligopeptide backbone. General references concerning the
coupling of proteins to primary amines or carboxylic acids
25 include Science, vol. 144, p. 1344 (1964); Erlanger et

al., Meth. in Immunology and Immunochemistry, Williams and Chase, eds, Academic, New York, p. 149 (1967); Kopple, Peptides and Amino Acids, Benjamin, New York (1966); Clin. Chem., vol. 22, p. 726 (1976); Immunochem., vol. 6, p. 53
5 (1969) and Lowe et al, Affinity Chromatography, Wiley, New York (1974).



Hapten ligands which themselves contain carboxylic acid functions, and which thereby can be coupled directly to the terminal -NH_2 group of the oligopeptide backbone,
10 include the iodothyronine hormones such as thyroxine and liothyronine, as well as other materials such as biotin, valproic acid, folic acid and certain prostaglandins. Representative synthetic routes for preparing carboxylic acid binding analogs of hapten ligands which themselves do
15 not contain an available carboxylic acid function, whereby such analogs can be coupled to the terminal -NH_2 group of the oligopeptide backbone by the aforementioned peptide condensation reactions or bifunctional coupling agent reactions, are given in U.S. Patent 4,331,808, which is
20 incorporated herein by reference. Thus, haptens such as carbamazepine, quinidine, digoxin, digitoxin, theophylline, phenobarbital, primidone, diphenylhydantoin, morphine, nicotine, androgens, estrogens, and progesterones may be bonded to the terminal NH_2 group of
25 the oligopeptide backbone. Compounds such as cocaine may

also be linked to the terminal -NH_2 group of the oligopeptide backbone. In addition, compounds which possess leaving groups may also be bonded to the terminal NH_2 group. For example, the trinitrophenyl or
5 dinitrophenyl group may be attached by the reaction of fluorotrinitrobenzene or fluorodinitrobenzene with the terminal NH_2 group of the oligopeptide.

As noted above, the ligand or receptor may be bonded to the oligopeptide backbone via a conventional
10 bifunctional spacer. The binding of a low molecular weight hapten, such as, e.g., the trinitrophenyl group, to the oligopeptide via a spacer can be advantageous in that the accessibility of the hapten for binding with its complement (antibody) may be increased. A suitable spacer
15 is, e.g., $\text{-(CH}_2\text{)}_m\text{-S-S-}$ (wherein m is 2 to 10, preferably 4 to 6). For example, the dinitrophenyl group may be bonded to the terminal -NH_2 group of the oligopeptide as shown below.



-16-



In the first step S-(2-thiopyridyl)-cysteamine prepared by the reaction of 2,2'-dithiopyridine with cysteamine in methanol (Chong et al, J. Biol. Chem., Vol. 256, pp. 5064-5070 (1981)), is reacted with dinitrofluorobenzene.

- 5 At the same time, the terminal -NH₂ group of the oligopeptide backbone is derivatized with Traut's reagent (2-iminothiolane) (Jue et al, Biochemistry, Vol. 17, pp. 5399-5406 (1978)). Then, the derivatized oligopeptide is coupled via the free sulfhydryl group to the pyridyldithio
- 10 group of the derivatized hapten.

- Other, bifunctional linkers which may be used to introduce spacers between the ligand or receptor and the oligopeptide backbone include those described in Haugland, Handbook of Fluorescent Probes and Research Chemicals,
- 15 Molecular Probes, Inc., Eugene, Oregon (1989) and Pierce Immunotechnology Catalog and Handbook, Pierce, Rockford, Il (1990), e.g., succinimidyl trans-4-(N-

maleimidylmethyl)-cyclohexane-1-carboxylate; succinimidyl
 4-(p-maleimidylphenyl)butyrate; succinimidyl
 6-((iodoacetyl)amino)hexanoate; succinimidyl 6-
 (6(((iodoacetyl)amino)hexanoyl)amino)hexanoate;
 5 succinimidyl 3-(2-pyridyldithio)propionate; and
 N-succinimidyl S-acetylthioacetate.

Thus, the oligopeptide and the ligand or receptor
 may be reacted with one of the above-mentioned molecules
 to introduce one of the following terminal functional
 10 groups:

polypeptide-SH

polypeptide-S-S-2-pyridyl

polypeptide-CCH₂I^O

polypeptide-maleimidyl


15 ligand-SH

ligand-S-S-2-pyridyl

ligand-CCH₂I^O

ligand-maleimidyl

The derivatized ligand or receptor may then be coupled
 20 with the derivatized oligopeptide as shown in the scheme
 below:



Alternatively, either the ligand or oligopeptide can be derivatized with N-succinimidyl S-acetylthioacetate resulting in a terminal S-acetyl group which can be deprotected with neutral NH_2OH to yield a free sulfhydryl group which can be coupled as shown above.

Thus, the ligand may be conveniently linked to the oligopeptide via a spacer which contains at least one of the following groups: thioether, disulfide, and thiosuccinimidyl.

- 10 Suitable polyoxoanions have the formula $-\text{OMO}_n$ and $-\text{MO}_n$, where M may be S or P and n is an integer of from 2 to 3, and include sulfate, sulfonate, sulfinatate, phosphate and phosphonate. These groups are easily introduced into amino acids and/or the amino acid residues of
- 15 polypeptides. For example, sulfate groups may be introduced into tyrosine by the reaction with sulfuric

acid at low temperatures (Reitz et al, J. Am. Chem. Soc.,
vol. 68, p. 1024 (1946) and Kohli et al, FEBS Lett., vol.
242, pp. 139-143 (1988)) and may be introduced into
tyrosine residues of polypeptides by the use of
5 concentrated sulfuric acid (Ondetti et al, J. Am. Chem.
Soc., Vol. 92, pp. 195-199 (1970). Sulfonate groups may
be introduced by either the conversion of the free
sulfhydryl groups of cysteine residues to S-sulfo-
derivatives by treatment with sodium tetrathionate, $\text{Na}_2\text{S}_4\text{O}_6$
10 (Dixon et al, Nature, vol. 188, p. 721 (1960)) or peptides
containing disulfide linkages can be reduced with
dithiothreitol followed by treatment with sodium
tetrathionate (Inglis et al, J. Biol. Chem., vol. 245, p.
112 (1970)). In addition, 3-sulfinol-L-alanine is
15 available from Aldrich and may be incorporated in the
oligopeptide backbone.

Phosphate groups may be introduced by a variety of
methods. Thus, tyrosine may be converted to the
O-phosphate derivative by reaction with P_2O_5 in H_3PO_4
20 (Rothberg et al, Proc. Natl. Acad. Sci. USA, vol. 75, pp.
4868-4872 (1978)) or serine may be converted to
phosphoserine by treatment with POCl_3 (Neuhaus et al,
Biochem. Prepn., vol. 6, p. 75 (1958)). Alternatively,
phosphothreonine is commercially available and may be used
25 as a starting amino acid for the production of the

polypeptide oligomer backbone. In addition, histidine can be derivatized to 7-phosphohistidine by reaction with phosphoamidate (Hulquist et al, Biochemistry, vol. 5, p. 22 (1966) and Fujitaki et al, Meth. Enzymol., vol. 107, 5 pp. 23-26 (1984)), and arginine can be derivatized to (ω -N-phosphoarginine by reaction with phosphorus oxychloride (Thiem et al, Bull. Soc. Chim. Biol., vol. 5, p. 322 (1962) and Fujitaki et al, Meth. Enzymol., vol. 107, pp. 23-26 (1984))).

10 Phosphonate moieties may be introduced into polypeptide oligomers by the reaction of the free sulfhydryl groups of cysteine residues with a phosphonic acid derivative, generated by the reaction of 3-aminopropylphosphonic acid with iodoacetic anhydride 15 (Meth. Enzymol., vol. 11, p. 532 (1967)).

In addition to the methods described above, phosphate groups may be introduced into polypeptides by the enzymatic phosphorylation of amino acid residues in polypeptides. For example, casein kinase of type II and 20 cAMP-dependent kinases phosphorylate serine, while casein kinases of type I phosphorylate threonine and, to a lesser extent, serine. Another group of kinases exhibits strict specificity for tyrosine (Corbin and Hardman, eds., Meth. Enzymol., vol. 99, Part F, Protein Kinases (1983)).

Similarly, amino acid residues in polypeptides may be enzymatically sulfonated. Thus, tyrosylprotein sulfotransferase catalyzes the sulfonation of proteins at tyrosine residues (Lee et al, J. Biol. Chem., vol. 258, p. 11326 (1983) and Huttner, Meth. Enzymol., vol. 107, pp. 200-233 (1984)).

It is preferred that the polyoxoanion is sulfonate.

The polyoxoanion groups may be present as free acids or may be partially or totally neutralized. In the salts formed by partial or total neutralization of the free acid form of the polyoxoanion, the cation may be any suitable ion such as a metal ion or an ammonium ion. Suitable metal ions are those that do not negate the water solubilizing effect of the polyoxoanion and include alkali metals such as lithium, sodium, potassium; alkaline earth metals such as magnesium, calcium, etc.; and transition metals such as, e.g., iron, copper, zinc, etc. Suitable ammonium ions include NH_4^+ and those in which one or more of the hydrogens have been substituted by an organic group, such as, e.g., tetramethylammonium.


The functional groups which serve to bond the fluorescent or chemiluminescent labels are suitably any of the reactive functional groups found in R of the naturally

occurring amino acids, e.g., -NH_2 in lysine (Lys);
 $\text{-NHC(NH}_2\text{)=NH}$ in arginine (Arg); -CONH_2 in asparagine (Asn)
and glutamine (Gln); -OH in serine (Ser), threonine (Thr),
and tyrosine (Tyr); and $\text{-CO}_2\text{H}$ in aspartic acid (Asp) and
5 glutamic acid (Glu). It is preferred that the functional
group be one of either $\text{NH}_2\text{-}$ and $\text{-CO}_2\text{H}$, so that one terminus
of the polypeptide backbone may also be conveniently used
as one of the sites for bonding the fluorescent or
chemiluminescent label.

10 It is to be understood that the -NH , groups in R of
e.g., Lys and the -NH_2 terminal position; the -CONH_2 groups
in R of Asn and Gln; the $\text{-NHC(NH}_2\text{)=NH}$ group in R of Arg;
and the imidazole group in R of His are sufficiently
dissimilar in reactivity to be classified as different
15 reactive groups for the purposes of the present invention.
In other words, the presence of an Asn or Arg residue in
the oligopeptide backbone does not prevent the terminal
 -NH_2 from serving as the unique functional group to bond
the ligand or receptor.

20 Specific examples of suitable chemiluminescent
labels are disclosed in German OLS No. 2,618,511, U.S.
Patent 4,331,808, and Haugland, Handbook of Fluorescent
Probes and Research Chemicals, Molecular Probes, Inc.,
Eugene, Oregon (1989) which are incorporated herein by


reference. Thus, suitable labels include luminol, isoluminol, pyrogallol, luciferin, and naphthalene-1,2-dicarboxylic acid hydrazide derivatives. These labels may be attached to the oligopeptide backbone by conventional methods. For example, a polypeptide may be labeled with a 7-aminonaphthalene-1,2-dicarboxylic acid hydrazide derivative by condensation of any $-CO_2H$ groups on the oligopeptide backbone with a compound of the formula:



as disclosed in U.S. Patent 4,331,808.

- Specific examples of suitable fluorescent labels are those disclosed in Blecka et al, "Immunoassays in Therapeutic Drug Monitoring," Clinics in Laboratory Medicine, vol. 7, pp. 357-370 (1987) and Haugland, Handbook of Fluorescent Probes and Research Chemicals, Molecular Probes, Inc., Eugene, Oregon (1989), which are incorporated herein by reference, and include, e.g., fluorescein, rhodamine, anthracene and fluoescamine. Again, the fluorescent labels may be attached to the

oligopeptide backbone by conventional methods. For example, fluorescein labels may be bonded to a polypeptide which contains a plurality of $-CO_2H$ groups by first treating the polypeptide with carbohydrazide, to obtain a
 5 carbohydrazide derivative, followed by treating the carbohydrazide derivative with fluorescein isothiocyanate (FITC). This procedure is shown schematically below:



Similarly, rhodamine X isothiocyanate and 2-anthraceneisothiocyanate may be coupled with the carbohydrazide derivative.

15 Further examples of labels include ruthenium complexes such as those described in U.S. Patent 4,745,076, incorporated herein by reference. These labels

are particularly suitable for time-resolved measurements (U.S. Patent 4,745,076) and electrogenerated chemiluminescent measurements (Ege et al, Analytical Chem., vol. 56, 2413 (1984) and Zhang et al, J. Phys. Chem., vol. 92, 5566 (1988)).

In regard to the number and distribution of the polyoxoanion groups and the groups for bonding the labels, it is preferred that there be at least two of each of these types of groups. It is especially preferred that
10 there be at least three of both the polyoxoanions and the groups for bonding the label. It is also preferred that the ratio of the number of polyoxoanions to the number of the functional groups for bonding the labels fall within the range of 1:4 to 4:1, more preferably 1:2 to 2:1. In
15 the interest of signal strength, it is preferred to keep the number of labels in the molecule relatively high. However, as the ratio of labels to anions in the molecule increases, the hydrophilicity and solubility decrease and the likelihood of signal quenching also increases. In
20 this regard, a suitable degree of labeling in the present molecules, and thus a suitable occurrence of groups for bonding the label, is on average about one label per 3 to 10 amino acid residues. A similar degree of substitution is suitable for the polyoxoanion.

It is to be understood that neither the functional groups for bonding the label nor the polyoxoanions need be spaced with strict regularity along the polypeptide backbone. Rather, it is only required that the labels be
5 spaced sufficiently far apart to avoid quenching and that the polyoxoanions be spaced such as to impart the desired hydrophilicity/solubility.

The amino residues contained in the backbone may be any of the naturally occurring α -amino acids represented
10 by the formula $H_2N-CH(R)-CO_2H$, such as glycine (Gly), alanine (Ala), valine (Val), leucine (Leu), isoleucine (Ile), methionine (Met), proline (Pro), phenylalanine (Phe), tryptophan (Trp) serine (Ser), threonine (Thr), cysteine (Cys), tyrosine (Tyr) asparagine (Asn), glutamine
15 (Gln), aspartic acid (Asp), glutamic acid (Glu), lysine (Lys), arginine (Arg), and histidine (His). The content and sequence of the amino acids are limited only by the need to satisfy the requirement of providing the three types of functional groups discussed above.

20 For example, if the terminal $-NH_2$ group of the polypeptide oligomer is to serve as the unique functional group for bonding to the ligand, then the polypeptide oligomer should not contain any amino acid residues which contain $-NH_2$ groups in R, such as, for example, lysine.

-27-

Similarly, when the terminal $\text{-CO}_2\text{H}$ group is to serve as the unique functional group, the backbone should contain no residues which contain $\text{-CO}_2\text{H}$ in R. On the other hand, a polypeptide which contains a plurality of aspartic acid
5 or glutamic acid residues will provide a number of $\text{-CO}_2\text{H}$ groups which can serve to bond the fluorescent or chemiluminescent label.

Thus, the oligopeptide backbone may be either a naturally occurring or synthesized oligopeptide. The
10 synthetic oligopeptide backbone may be synthesized by any conventional procedure. In particular, solid state synthesis utilizing any of the commercially available instruments, such as that produced by Applied Biosystems of Foster City, CA, is suitable. A discussion of the
15 solid state synthesis of oligopeptides is provided in U.S. Patent 3,531,258, which is incorporated herein by reference.

Alternatively, the oligopeptide may be synthesized by recombinant DNA technology. That is, the gene encoding
20 for the desired oligopeptide may be synthesized or isolated and then inserted into a suitable cloning vector which is used to transform a suitable host. The production of oligopeptides by recombinant DNA technology is discussed in U.S. Patents 4,704,362, 4,652,525


4,431,740, 4,440,859, and 4,342,832, which are incorporated herein by reference.

The present conjugates may be assembled by first introducing the polyoxoanion groups into the oligopeptide backbone, followed by bonding the ligand or receptor to the unique functional group and then attaching the labels to the remaining functional groups. Alternatively, the labels may be attached to the backbone before the ligand or receptor is attached or the polyoxoanions may be introduced after either of the other two steps. Thus, in one embodiment, the present invention relates to an intermediate ligand compound which is an oligopeptide, bonded to a ligand or receptor, in which at least one of the amino acid residues has been converted to a polyoxoanion derivative and a plurality of the amino acid residues contain a functional group for bonding a fluorescent or chemiluminescent label. In another embodiment, the present invention relates to an intermediate labeled compound which is an oligopeptide, in which at least one of the amino acid residues has been converted to a polyoxoanion derivative, a plurality of the amino acid residues are linked to a fluorescent or chemiluminescent label, and having a unique functional group for bonding a ligand or receptor.

These intermediate compounds may be represented by the general formulae




5 wherein L and n are as defined above and at least one of the amino acid residues contains a polyoxoanion of phosphorus or sulfur and a plurality of the amino acid residues have a functional group for bonding a chemiluminescent or fluorescent label; or



wherein n is as defined above and at least one of the amino acid residues contains a polyoxoanion of phosphorus or sulfur and a plurality of the amino acid residues are bonded to a chemiluminescent or fluorescent label.

15 Examples of suitable oligopeptide backbones include the insulin A-chain and fragments of cholecystokinin-pancreozymin which may be sulfated as disclosed in Ondetti et al, J. Am. Chem. Soc., Vol. 92, pp. 195-199 (1970) some of which are shown below.



A preferred embodiment of the oligopeptide backbone of present conjugates is represented by insulin A-chain or

5 a polypeptide having substantially the same number of amino acid residues and molecular weight as insulin A-chain, with one terminal amino or carboxy group for covalent attachment of a ligand or receptor molecule, at least two functional groups for covalent attachment of a

10 label at a distance from the ligand or receptor site sufficient to minimize quenching of the labels by the ligand or receptor, and at least two functional groups which may be converted to a polyoxoanion of phosphorus or sulfur. Such oligopeptides include insulin A-chain;

15 oligopeptides having substantially the same number of amino acid residues as insulin A-chain, at least two cysteine residues and at least one amino acid residue having a $-\text{CO}_2\text{H}$ group in R, such as Glu or Asp, but only one $-\text{NH}_2$ group; and oligopeptides having substantially the

20 same number of amino acid residues as insulin A-chain, at least two cysteine residues, and at least one amino acid

residue having a $-NH_2$ group in R, such as Lys, but only one $-CO_2H$ group. Examples of such peptides may be represented by the formula




- 5 in which n is about 21, at least two of the amino acid residues are cysteine, at least one of the amino acid residues has a $-CO_2H$ group in R, such as Glu or Asp, and none of the remaining amino acid residues have $-NH_2$ groups in R, such as Lys; or n is defined as above, at least two
10 of the amino acid residues are cysteine, at least one of the amino acid residues has a $-NH_2$ group in R, such as Lys, and none of the remaining amino acid residues contain a $-CO_2H$ group in R, such as Glu or Asp.

The A-chain of insulin represents an ideal backbone
15 for the synthesis of a sulfonated oligomeric carrier molecule. The A-chain of insulin is a hydrophilic peptide containing 21 amino acid residues and has a molecular weight of approximately 2500 daltons.

Four of the amino acids are cysteine residues with a
20 free sulfhydryl group which can easily be derivatized to S-sulfonates, J. Am. Chem. Soc., vol. 88, 5625-5635 (1966). Two carboxyl groups from glutamate residues 4 and

17 and one terminal carboxyl group from an asparagine residue provide three sites for attachment of fluorescent or chemiluminescent molecules. The insulin A-chain contains one terminal amino group for covalent attachment
5 of a single ligand or receptor molecule. Since the sites for attachment of the fluorescent or chemiluminescent labels are 4, 17, and 21 amino acid residues away from the ligand or receptor attachment site, the possibility of quenching of the labels by the ligand or receptor is
10 minimized.

Thus, insulin A-chain in the tetra-S-sulfonate form may be easily coupled to an antigen, such as the dinitrophenyl group, via the terminal $-NH_2$ group by reaction with 1-fluoro-2,4-dinitrobenzene, to obtain
15 dinitrophenyl-insulin A-chain (DNP-insulin A-chain). Derivatization of the carboxyl groups of DNP-insulin A-chain with carbohydrazide, followed by reaction with fluorescein isothiocyanate (FITC) gives DNP-insulin A-chain-fluorescein (DNP-Ins-Fl), the structure of which
20 is shown below:



The ligand-insulin A-chain-fluorescein conjugates (Ligand-Ins-Fl) are ideally suited for use in assays for the free ligand. For example, the amount of non-specific binding of DNP-Ins-Fl to immobilized anti-DNP IgG (antibody #51) is only about one-third that of a conjugate in which a dinitrophenyl group is linked to fluorescein via a lysine residue, DNP-Lys-fluorescein (DNP-Lys-Fl) (see Table II). Further, the amount of DNP-Ins-Fl specifically bound to antibody #51 is about 1.7-fold higher than that of DNP-Lys-Fl.

In Figure 3 the binding of the different DNP conjugates to anti-DNP-Sepharose at a 3:1 molar ratio of conjugate to immobilized antibody is shown. After extensive washing of the matrix, a 1.6-fold difference in binding of DNP-Lys-Fl (62.7%) versus DNP-Ins-Fl (38.8%)

was observed. In an attempt to identify the factors contributing to this significant difference, the binding of DNP-Ins without attached fluorescein molecules was also determined. The data in Table II reveal an almost
5 identical extent of binding of DNP-Ins and DNP-Ins-F1 to anti-DNP-Sepharose, suggesting that the increased number of fluorophores in the DNP-Ins-F1 conjugate compared to the DNP-Lys-F1 conjugate is not a critical factor.

In order to determine whether the differences in the
10 extent of binding shown in Figure 3 are a result of a different affinity of the immobilized anti-DNP antibody for both conjugates, DNP-Ins-F1 and DNP-Lys-F1, the displacement of the different DNP conjugates from the antigen binding sites of immobilized anti-DNP antibody #51
15 by DNP-lysine was determined. The analyses revealed that under comparable conditions, the displacement of DNP-Ins-F1 is much more efficient than that of DNP-Lys-F1 (see Table III). However, the data in Table III also demonstrate that the displacement of DNP-Ins-F1 is two-
20 fold more efficient than that of DNP-Ins, whereas the extent of binding of DNP-Ins-F1 and DNP-Ins was almost identical (Table II). Therefore, a difference in affinity cannot be the only factor responsible for the different binding efficiencies of the DNP conjugates to antiDNP-
25 Sepharose.

Table II compares the extent of binding of the different DNP conjugates to immobilized anti-DNP IgG (antibody #51) with that to immobilized non-immune IgG. The data show that the extent of non-specific binding of DNP-Lys-F1 to immobilized non-immune IgG is three-fold higher than that of DNP-Ins-F1. After subtraction of non-specific binding to non-immune IgG-Sepharose from binding to anti-DNP-Sepharose, a fundamentally different binding pattern becomes evident. The amount of DNP-Ins-F1 that specifically is bound to the antigen binding sites of immobilized antibody #51 (0.7 mole conjugate/mole antibody) is 1.7-fold higher than that of DNP-Lys-F1 (0.4 mole conjugate/mole antibody). These results show that the different extent of binding of DNP-Lys-F1 and DNP-Ins-F1 to anti-DNP-Sepharose given in Figures 3 is a consequence of the high nonspecific binding of DNP-Lys-F1.

In another aspect, the present invention relates to kits which contain the present conjugates and/or intermediate compounds. Such kits may take any form that is suitable for the employment of the present conjugates and/or intermediate compounds in an assay for a ligand and/or receptor. In particular, the present kits may contain a vial or other container which may contain the present conjugate and/or intermediate compound in solution

or powdered form. If the present conjugate and/or intermediate compound is in solution form, the solution may also contain other materials, such as, e.g., a preservative, buffer, salt, etc. The kit may also contain
5 a standard sample of either the ligand or receptor of the present conjugate or the binding complement of the ligand or receptor. Such standard samples may be in the form of a solution of known concentration and/or activity or a sample of known weight and/or activity. In addition, the
10 present kits may contain samples in which the binding complement of the ligand or receptor of the conjugate is immobilized on a solid support, such as a glass slide, glass beads, gelatin beads, polymer matrix, etc.

The present conjugates and kits are useful for
15 assaying ligands or receptors and/or the binding complements of the ligands or receptors, in specific binding assays in which the displacement or competitive binding of the present conjugates are used to determine the amount or concentration of the free ligand or receptor
20 (or complement thereof) present in a sample. The sample may be of biological or nonbiological origin and may be originally obtained in gaseous, liquid or solid form.

Other features of the invention will become apparent in the course of the following descriptions of exemplary

embodiments which are given for illustration of the invention and are not intended to be limiting thereof.

General Methods

Fluorescence intensity was measured on a SLM 8000
5 fluorimeter at 490 nm excitation and 519 nm emission.
Individual fluorescence standard curves with known
concentrations of the different DNP-fluorescein conjugates
were used to calculate the amounts of DNP-conjugates from
fluorescence units. Antibodies were coupled to tressyl
10 chloride-activated Sepharose 4B (Sigma) at a concentration
of 200 ug antibody/100 mg gel according to the
manufacturer's recommendation. The amount of immobilized
antibody was determined using a Coomassie blue assay as
described in Ahmad et al, Anal. Biochem., vol. 148, pp.
15 533-541 (1985). Peptide and protein determinations in
fluid-phase were performed by UV absorbance at 280 nm, by
the bicinchoninic acid method (Smith et al, Anal.
Biochem., vol. 150, pp. 76-85 (1985)) using the Pierce BCA
Protein Assay Reagent (Pierce), and by the FolinCiocalteu
20 method (Lowry et al, J. Biol. Chem., vol. 193, pp. 265
(1951)). The A-chain of bovine insulin (oxidized to the
S-sulfonate form) (Sigma), the DNP-Ins conjugate, and the
DNPIns-Flu conjugate were radiolabeled with Na¹²⁵I
(Amersham, Arlington Heights, IL) using immobilized

chloramine-T (IodoBeads; Pierce, Rockford, IL) (Petrella et al, J. Immunol. Methods, vol. 104, pp. 159-172 (1987)).

Synthesis of the DNP-Insulin A-Chain-Fluorescein Conjugate

The synthesis was performed in three steps: 1)
5 coupling of 1-fluoro-2,4-dinitrobenzene (FDNB) to the
terminal amino group of the insulin A-chain oxidized to
the tetra-S-sulfonate form, 2) derivatization of the
carboxyl groups with carbohydrazide, and 3) coupling of
fluorescein isothiocyanate (FITC) to the hydrazide
10 derivative of DNP-insulin A-chain.

Step 1. Insulin A-chain in the tetra-S-sulfonate
form (Sigma, St. Louis, MO) (25 mg; approximately 8
 μ moles), mixed with ^{125}I -labeled S-sulfonated insulin
A-chain (26 nmole, 5×10^7 cpm), was incubated with 50 μ mol
15 (9.3 mg) of FDNB (Aldrich, Milwaukee, WI) in a total
volume of 6.4 mL containing 100 mM NaHCO_3 and 19% (v/v)
ethanol. After two hours of incubation at 37°C, the
reaction mixture was subjected to Sephadex G-10 size
exclusion chromatography to remove noncoupled FDNB
20 molecules. The peptide fractions, eluted with 19% (v/v)
ethanol in H_2O , were lyophilized and stored at -20°C. The
ratio of coupled FDNB residues per insulin A-chain was
calculated from the specific radioactivity of the peptide

and the UV absorbance at 365 nm using an extinction coefficient of $1.6 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ (Hirs, Meth. Enzymol., XI, pp. 548-555 (1967)). Preparations containing less than one DNP residue per insulin A-chain were subjected to
5 a second derivatization with freshly added FDNB as described above.

Step 2. DNP-derivatized insulin A-chain (tetra-S-sulfonate form) (DNP-Ins) (3.5 mg; approximately $1.2 \mu\text{mol}$) was dissolved in 800 μl of 10 mM sodium phosphate-150 mM
10 NaCl, pH 7.4. After the addition of 165 mg (1.8 mmol) of carbohydrazide (Aldrich) and 20 mg (0.1 mmol) of 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide (EDC) (Sigma), the pH was adjusted to pH 5.0 with hydrochloric acid. The reaction mixture (total volume 950 μl) was incubated at
15 room temperature for 18 hours (the pH was maintained at 5.0) and then subjected to Sephadex G-10 size exclusion chromatography. The peptide fraction, eluted with 19% (v/v) ethanol in H_2O , was lyophilized and stored at -20°C .

Step 3. Carbohydrazide-derivatized DNP-insulin
20 A-chain (tetra-S-sulfonate form) (0.35 mg; approximately 0.1 nmol) was dissolved in 950 μl of 50 mM NaHCO_3 containing 19% (v/v) ethanol. FITC (Aldrich) (16 mg; $41 \mu\text{mol}$) was added, and after 18 hours of incubation at room temperature in the dark, the reaction mixture was

subjected to Sephadex G-10 size exclusion chromatography. The peptide fraction, eluted with 19% (v/v) ethanol in H₂O, was lyophilized, redissolved in 50 mM NaHCO₃ containing 19% (v/v) ethanol, and rechromatographed on

5 Sephadex G-10 to remove traces of any non-covalently bound fluorophores. The number of covalently attached fluorescein molecules was spectrophotometrically determined at 494 nm using an extinction coefficient of $7.6 \times 10^4 \text{ M}^{-1}\text{cm}^{-1}$ (Wilderspin et al, Anal. Biochem., vol.

10 132, pp. 449-455 (1983)).

Spectrophotometric analyses of the purified DNP-insulinA-chain-fluorescein conjugate (DNP-Ins-F1) confirmed the theoretical ratio of one DNP residue and three fluorescein residues per insulin A-chain molecule as


15 shown below in Table I.

Table I. Molar ratio of covalently attached FDNB and FITC molecules per insulin A-chain.

20	Derivatization of A-chain insulin A-chain with:	Ratio: modifying residue/insulin	
		Expected	Found
	FDNB	1.0	1.1 ± 0.1
25	FITC	3.0	3.2 ± 0.1

Synthesis of the DNP-Lysine -Fluorescein Conjugate

ϵ -DNP-L-lysine hydrochloride (Research Organics Inc., Cleveland, Ohio 44125) (10 mg; 0.029 mmol) was dissolved in 1.0 ml of 0.5 M NaHCO₃ and adjusted to pH 9.3 with NaOH. Fluorescein isothiocyanate (Aldrich) (10 mg; 0.026 mmol) was added, and the reaction mixture (total volume 1.2 ml) was incubated in the dark for 30 min at room temperature. Purification of the synthesized DNP-lysine-fluorescein conjugate (DNP-Lys-Fl) was performed by reverse phase HPLC on an Ultrasphere ODS C-18 column (Beckman, CA) at a flow rate of 0.75 ml/min using a linear gradient from 5% (v/v) methanol (in H₂O) to 100% methanol (within 15 min). The DNP-Lys-Fl conjugate fractions from several chromatographic runs were combined, lyophilized, and stored at -20°C. Rechromatography of an aliquot of the combined fractions under identical conditions demonstrated 95% purity of the DNP-Lys-Fl conjugate. The structure of DNP-Lys-Fl is shown below.



Determination of Fluorescence Lifetime

The fluorescence lifetime of the DNP-fluorescein conjugates was measured on an ISS Greg-200 variable frequency phase fluorimeter as described previously (Lakowicz et al, Biophys. J., vol. 46, pp. 463-477 (1984) and Gratton et al, Biophys. J., vol. 46, pp. 479-486 (1984)) using 1-chloro-bis-(phenylethynyl)anthracene (Aldrich) in ethanol (3.791 ns) as a reference (Thompson et al, Anal. Chem., vol. 60, pp. 670-674 (1988)). The resulting phase and modulation data for DNP-Ins-FL were fit to one, two, and three emissive components using a Simplex algorithm supplied by ISS. The results are shown in Figure 1. The data fitted best ($\chi^2=1.26$) to two components: the first comprises about two thirds of the mixture and exhibited a lifetime corresponding closely to that of fluorescein (4.091 ± 0.177 ns; $64 \pm 7\%$), and a shorter lifetime component comprising about one third of the mixture (2.068 ± 0.150 ns; 36%). Fitting these data to one component gave an unacceptable χ^2 ($\chi^2 = 33$) and non-random differences between the data and calculated values. When the data were fit to a three component model, the third component was weak (<5%), poorly defined (1.1 ± 3.2 ns), and gave no improvement in χ^2 . Therefore, a third component appears to be unnecessary to fit the data. Since the DNP moiety has the ability to quench the

fluorescein emission, the data are best explained by two fluorescein residues being remote from and unquenched by the DNP group with the third fluorescein residue being closer and subject to quenching. In accordance with this
5 explanation is the observation that the fluorescence of the DNP-lysine-fluorescein derivative was substantially quenched as shown by its reduced quantum efficiency and subnanosecond fluorescence lifetime.

In addition, the relationship between the
10 fluorescence intensity and concentration for DNP-Ins-Fl and DNP-Lys-Fl is given in Figure 2.

Specific and Nonspecific Binding of DNP-Conjugates

Specific binding of the DNP-conjugates was determined with the murine monoclonal anti-DNP antibody
15 #51 (Stanley et al, J. Immunol. Methods, vol. 64, pp. 119-132 (1983) and Steward et al, J. Immunol. Methods, vol. 78, pp. 173-190 (1985)) coupled to tressyl chloride-activated Sepharose 4B (anti-DNP-Sepharose). The monoclonal IgG1 antibody #51 (purified from ascites) was
20 kindly provided by Dr. M.W. Steward (Dept. of Medical Microbiology, London School of Hygiene and Tropical Medicine, London, U.K.). Nonspecific binding was measured with Sepharose 4B-immobilized non-immune mouse IgG

purified from mouse serum (Jackson ImmunoResearch Laboratories, West Grove, PA) (IgG-Sepharose). After removal of buffer by vacuum filtration on a scintered glass funnel, 50 mg aliquots of Sepharose 4B-antibody

5 matrices were transferred to plastic conical vials and incubated with ^{125}I -labeled DNPinsulin A-chain-fluorescein conjugate (1.6×10^6 cpm/nmole), or ^{125}I -labeled DNP-insulin A-chain conjugate without attached fluorescein residues (3.3×10^6 cpm/nmole), or non-radiolabeled DNP-

10 Lys-F1 at a 3 to 1 molar ratio of DNP-derivative to immobilized IgG in 10 mM phosphate/150 mM NaCl, pH 7.6, overnight at 4°C on a rocking platform. Unbound conjugate was recovered by poking a hole in the bottom of the plastic vial with a 30 gauge needle, loading the vial on

15 the top of a glass test tube, and centrifuging. Subsequent washing steps were performed with 150 μl of 10 mM phosphate/150 mM NaCl/0.1% (v/v) Triton X-100, pH 7.6. Aliquots of the solutions collected in the bottom of the test tubes after centrifugation were assayed for

20 radioactivity or fluorescence. The results are shown in Table II and Figure 3.

Table II. Specific and nonspecific binding of DNP-conjugates.

5	DNP-Conjugate	DNP-Conjugate Applied	Ratio of Applied DNP-Conjugate to Immobilized IgG	DNP-Conjugate Bound	Ratio of Bound DNP-Conjugate to Immobilized IgG
		(nmoles)	(moles/moles)	(nmoles)	(moles/moles)
10					
15	I. Binding to anti-DNP IgG1				
	DNP-Lys-F1	0.59	3.0	0.37±0.02	1.9
	DNP-Ins-F1	0.29	3.0	0.11±0.03	1.2
	DNP-Ins	0.29	3.0	0.11±0.01	1.2
20	II. Binding to non-immune IgG				
	DNP-Lys-F1	2.60	3.0	1.27±0.07	1.5
	DNP-Ins-F1	2.53	3.0	0.44±0.16	0.5
25					

Specificity and Effectivity of Displacement of DNP-Conjugates

Aliquots of anti-DNP-Sepharose (50 mg) were incubated with ^{125}I -labeled DNP-insulin A-chain, or ^{125}I -labeled DNPinsulin A-chain-fluorescein, or DNP-lysine-fluorescein and washed with 10 mM phosphate/150 mM

NaCl/0.1% (v/v) Triton X100, pH 7.6, as described above in Specific and Nonspecific Binding of DNP-Conjugates. When the wash solutions showed background levels of radioactivity or fluorescence, 150 μ l of 10 mM

5 phosphate/150 mM NaCl/0.1% (v/v) Triton X-100, pH 7.6, containing the compound to be tested for displacement of antibody-bound DNP-conjugate was added to the anti-DNP-Sepharose aliquots, and centrifuged as described.

Displacement was determined by assaying aliquots of the
10 solutions collected in the bottom of the test tubes after centrifugation for radioactivity or fluorescence. The results are shown in Table III.

15 Table III. Displacement of DNP-conjugates from immobilized monoclonal anti-DNP antibody by DNP-lysine.

20	Antibody-Bound DNP-Conjugate	Ratio of Free DNP-Lysine to Antibody-Bound DNP-Conjugate	Displacement of Antibody-Bound DNP-Conjugate by DNP-Lysine
		[mole/mole]	[pmole \pm S.D.]
	DNP-Lys-F1	2.0	0.45 \pm 0.14
	DNP-Ins-F1	3.4	3.20 \pm 0.70
25	DNP-Ins	3.4	1.56 \pm 0.01

Obviously, numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore to be understood that within the scope of the appended claims, the invention may be
5 practiced otherwise than as specifically described herein.

CLAIMS

1. A ligand-label conjugate, which is an oligopeptide of 5 to 100 amino acid residues, which is bonded to a ligand or receptor, wherein at least one of said amino acid residues contains a polyoxoanion of sulfur
5 or phosphorus and a plurality of said amino acid residues are linked to a chemiluminescent or fluorescent label.

2. The conjugate of Claim 1, wherein the number of amino acid residues in said oligopeptide is from 10 to 50.

3. The conjugate of Claim 1, wherein the number of
10 amino acid residues in said oligopeptide is from 15 to 25.

4. The conjugate of Claim 1, wherein said label is selected from the group consisting of luminol, isoluminol, pyrogallol, luciferin, naphthalene-1,2-dicarboxylic acid hydrazide derivatives, fluorescein, rhodamine, anthracene,
15 fluorescamine, and ruthenium complexes.


5. The conjugate of Claim 1, wherein said polyoxoanion of sulfur or phosphorus is selected from the group consisting of sulfate, sulfonate, sulfinde, phosphate and phosphonate.

6. The conjugate of Claim 1, wherein said polyoxoanion of sulfur or phosphorus is sulfonate.

7. The conjugate of Claim 1, wherein said oligopeptide has substantially the same amino acid residues and molecular weight as insulin A-chain, said amino acid residues containing a polyoxoanion of sulfur or phosphorus are S-sulfonate-Cys, said amino acid residues linked to said labels are Glu or Asp and the CO₂H-terminal amino acid residue, and said ligand is linked to the H₂N-terminal amino acid residue.

8. The conjugate of Claim 1, wherein said oligopeptide has the same amino acid sequence as insulin A-chain.

9. The conjugate of Claim 1, which has the formula:



10. The conjugate of Claim 1, wherein said ligand or receptor is bonded to said oligopeptide via a spacer group.

11. The conjugate of Claim 10, wherein said spacer group contains a thioether group, a disulfide group, or a thiosuccinimidyl group.

12. A ligand compound which is an oligopeptide of 5 to 100 amino acid residues which is bonded to a ligand or receptor, wherein at least one of said amino acid residues contains a polyoxoanion of phosphorus or sulfur and a plurality of said amino acid residues contain a functional group for bonding a chemiluminescent or fluorescent label.

13. The ligand compound of Claim 12, wherein the number of amino acid residues in said oligopeptide is from 10 to 50.

14. The ligand compound of Claim 12, wherein the number of amino acid residues in said oligopeptide is from 15 to 25.

15. The ligand compound of Claim 12, wherein said polyoxoanion of sulfur or phosphorus is selected from the


group consisting of sulfat , sulfonate, sulfinat,
phosphat and phosphonat .

16. The ligand compound of Claim 12, wherein said
polyoxoanion of sulfur or phosphorus is sulfonate.

5 17. The ligand compound of Claim 12, wherein said
oligopeptide has substantially the same amino acid
residues and molecular weight as insulin A-chain, said
amino acid residues containing a polyoxoanion of sulfur or
phosphorus are S-sulfonate-Cys, said amino acid residues
10 linked to said labels are Glu or Asp and the CO₂H-terminal
residue, and said ligand is linked to said H₂N-terminal
amino acid residue.

18. The ligand compound of Claim 12, wherein said
oligopeptide has the same amino acid sequence as insulin
15 Achain.

19. The ligand compound of Claim 12, which has the
formula:



20. The ligand compound of Claim 12, wherein said functional group for bonding a chemiluminescent or fluorescent label is -NH_2 or $\text{-CO}_2\text{H}$.

21. The ligand compound of Claim 12, wherein said
5 ligand or receptor is bonded to said oligopeptide via a spacer group.

22. The ligand compound of Claim 21, wherein said spacer group contains a thioether group, a disulfide group, or a thiosuccinimidyl group.

10 23. A labeled compound which is an oligopeptide of 5 to 100 amino acid residues in which at least one of said amino acid residues contains a polyoxoanion and a plurality of said amino acid residues are bonded to a chemiluminescent or fluorescent label and one of said
15 amino acid residues contains a unique functional group for bonding a ligand or receptor.

24. The labeled compound of Claim 23, wherein the number of amino acid residues in said oligopeptide is from 10 to 50.

25. The labeled compound of Claim 23, wherein the number of amino acid residues in said oligopeptide is from 15 to 25.

26. The labeled compound of Claim 23, wherein said
5 chemiluminescent label is selected from the group consisting of luminol, isoluminol, pyrogallol, luciferin, naphthalene-1,2-dicarboxylic acid hydrazide derivatives, fluorescein, rhodamine, anthracene, fluorescamine, and ruthenium complexes.

10 27. The labeled compound of Claim 23, wherein said polyoxoanion of sulfur or phosphorus is selected from the group consisting of sulfate, sulfonate, sulfinatate, phosphate and phosphonate.

28. The labeled compound of Claim 23, wherein said
15 polyoxoanion of sulfur or phosphorus is sulfonate.


29. The labeled compound of Claim 23, wherein said unique functional group for bonding a ligand or receptor is $-NH_2$ or $-CO_2H$.

30. The labeled compound of Claim 23, wherein said
20 oligopeptide has substantially the same amino acid residues and molecular weight as insulin A-chain, said

amino acid residues containing a polyoxoanion of sulfur or phosphorus are S-sulfonated -Cys, said amino acid residues linked to said labels are Glu or Asp and the CO₂H-terminal residue, and said ligand is linked to the H₂N-terminal amino acid residue.

31. The labeled compound of Claim 23, wherein said oligopeptide has the same amino acid sequence as insulin Achain.

32. The labeled compound of Claim 23, which has the formula:



33. A kit, comprising (i) a ligand-label conjugate, which is an oligopeptide of 5 to 100 amino acid residues, which is bonded to a ligand or a receptor, wherein at

least one of said amino acid residues contains a polyoxoanion of sulfur or phosphorus and a plurality of said amino acid residues are linked to a chemiluminescent or fluorescent label, and (ii) a binding complement of
5 said ligand or receptor.

34. The kit of Claim 33, wherein the number of amino acid residues in said oligopeptide is from 10 to 50.

35. The kit of Claim 33, wherein the number of
10 amino acid residues in said oligopeptide is from 15 to 25..

36. The kit of Claim 33, wherein said chemiluminescent label is selected from the group consisting of luminol, isoluminol, pyrogallol, luciferin,
15 naphthalene-1,2-dicarboxylic acid hydrazide derivatives, fluorescein, rhodamine, anthracene, fluorescamine, and ruthenium complexes.


37. The kit of Claim 33, wherein said polyoxoanion of sulfur or phosphorus is selected from the group
20 consisting of sulfate, sulfonate, sulfinic acid, phosphate and phosphonate.

38. The kit of Claim 33, wherein said polyoxoanion of sulfur or phosphorus is sulfonate.

39. The kit of Claim 33, wherein said oligopeptide has substantially the same amino acid residues and molecular weight as insulin A-chain, said amino acid residues containing a polyoxoanion of sulfur or phosphorus are S-sulfonate-Cys, said amino acid residues linked to the labels are Glu or Asp and the CO₂H-terminal residue, and said ligand is linked to the H₂N-terminal amino acid residue.

40. The kit of Claim 33, wherein said oligopeptide
10 has the same amino acid sequence as insulin A-chain.

41. The kit of Claim 33, wherein said ligand-label conjugate has the formula:



42. The kit of Claim 33, further comprising a sample of said ligand or receptor.

43. The kit of Claim 42, wherein said sample of said ligand or receptor contains a known amount of said
5 ligand or receptor.

44. The kit of Claim 33, wherein said binding complement of said ligand or receptor is present in the form of a sample of a known amount.

45. The kit of Claim 33, wherein said ligand or
10 receptor is bonded to said oligopeptide via a spacer group.

46. The kit of Claim 45, wherein said spacer group contains a thioether group, a disulfide group, or a thiosuccinimidyl group.

1/3




FIG. 1

SUBSTITUTE SHEET

2/3




FIG. 2

SUBSTITUTE SHEET

3/3




FIG. 3

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US91/02212

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all.) According to International Patent Classification (IPC) or to both National Classification and IPC IPC(5): C07K 7/40; G01N 33/532, 33/533 U.S. CL: 435/964,968; 436/543,544,546,547,800,819,822,823; 530/303		
II. FIELDS SEARCHED Minimum Documentation Searched ¹		
Classification System	Classification Symbols	
U.S. CL.:	435/964,968; 436/543,544,546,547,800,819,822,823; 530/303	
Documentation Searched other than Minimum Documentation, to the Extent that such Documents are Included in the Fields Searched ²		
CASONLINE SEARCH: AMINO ACID SEQUENCE OF CLAIM 9 STRUCTURE; APS SEARCH: TERMS "INSULIN A", "SULFONAT?", "SULFINAT?", "SULFAT?", "PHOSPHAT?", "PHOSPHONAT?"		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ³		
Category ⁴	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X Y	US, 3, 4, 014, 861 (Geiger et al.) 29 March 1977, see column 2, formula (IV) and lines 7-11 and 36.	12-22 1-11, 23-46
Y	US, A, 4, 166, 105 (Hirschfeld) 28 August 1979, see column 1, line 42-column 2, line 8; column 2, lines 14-19; column 3, line 64-column 3, line 16; claim 1.	1-46
Y	US, A, 4, 610, 868 (Fountain et al.) 09 September 1986, see column 4, lines 47-50.	1-46
Y	US, A, 4, 604, 364 (Kosak) 05 August 1986, see claim 4.	1-46
Y	US, A, 4, 645, 646 (Gadow et al.) 24 February 1987, see column 2, lines 53-55 and claim 8.	1-46
A	US, A, 4, 430, 266 (Frank) 07 February 1984, see column 1, lines 63-67.	1-46
<p>⁴ Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search 13 JUNE 1991		Date of Mailing of this International Search Report 15 JUL 1991
International Searching Authority ISA/US		Signature of Authorized Officer MARY E. CEPERLEY

